

CUTANEOUS SQUAMOUS CELL CARCINOMA PROFILE

Procedure: Biopsy, punch
Biopsy, shave
Biopsy, other: _____
Excision, ellipse
Excision, wide
Excision, other: _____
Lymphadenectomy, sentinel node(s)
Lymphadenectomy, regional nodes (specify): ____
Other (specify): _____ -
Not specified

Tumor site: _____
Not specified

Tumor size (greatest dimension): ____ cm

Histologic type: (select all that apply)
Squamous cell carcinoma
Acantholytic SCC
Spindle cell (sarcomatoid) SCC
Verrucous SCC
Pseudovascular SCC
Adenosquamous carcinoma
Squamous cell carcinoma, type not otherwise specified
Other: _____

Histologic grade: GX: Cannot be assessed
G1: Well differentiated
G2: Moderately differentiated
G3: Poorly differentiated
G4: Undifferentiated

Maximum tumor thickness (Breslow): Not applicable
____ mm
At least ____ mm

Anatomic (Clark's) level: Not applicable
I: Carcinoma in situ
II: Carcinoma present in bot does not fill and expand papillary dermis
III: Carcinoma fills and expands papillary dermis
IV: Carcinoma invades reticular dermis
V: Carcinoma invades subcutaneum

Lymph-vascular invasion: Not identified
Present
Indeterminate

Perineural invasion: Not identified
Present
Indeterminate

Margins

Peripheral Margins

☐ Cannot be assessed
☐ Uninvolved by invasive carcinoma
Distance of invasive carcinoma from closest peripheral margin: ____ mm
Specify location(s), if possible: _____
☐ Involved by invasive carcinoma
Specify location(s), if possible: _____
☐ Uninvolved by carcinoma in situ
Distance of carcinoma in situ from closest peripheral margin: ____ mm
Specify location(s), if possible: _____
☐ Involved by carcinoma in situ
Specify location(s), if possible: _____

Deep Margin

☐ Cannot be assessed
☐ Uninvolved by invasive carcinoma
Distance of invasive carcinoma from margin: ____ mm
Specify location(s), if possible: _____
☐ Involved by invasive carcinoma
Specify location(s), if possible: _____

Pathologic Staging (pTNM)**Primary Tumor (pT)**

- ☐ pTX: Primary tumor cannot be assessed
- ☐ pT0: No evidence of primary tumor
- ☐ pTis: Carcinoma in situ
- ☐ pT1: Tumor 2 cm or less in greatest dimension with fewer than two high-risk features
- ☐ pT2: Tumor greater than 2 cm in greatest dimension with or without one additional high-risk feature, or any size with two or more high-risk features
- ☐ pT3: Tumor with invasion of maxilla, mandible, orbit, or temporal bone
- ☐ pT4: Tumor with invasion of skeleton (axial or appendicular) or perineural invasion of skull base

Regional Lymph Nodes (pN)

- ☐ pNX: Regional lymph nodes cannot be assessed
- ☐ pN0: No regional lymph node metastasis
- ☐ pN1: Metastasis in a single ipsilateral lymph node, 3 cm or less in greatest dimension
- ☐ pN2: Metastasis in a single ipsilateral lymph node, more than 3 cm but not more than 6 cm in greatest dimension; or in multiple ipsilateral lymph nodes, none more than 6 cm in greatest dimension; or in bilateral or contralateral lymph nodes, none more than 6 cm in greatest dimension
- ☐ pN2a: Metastasis in a single ipsilateral lymph node, more than 3 cm but not more than 6 cm in greatest dimension
- ☐ pN2b: Metastasis in multiple ipsilateral lymph nodes, none more than 6 cm in greatest dimension
- ☐ pN2c: Metastasis in bilateral or contralateral lymph nodes, none more than 6 cm in greatest dimension.
- ☐ pN3: Metastasis in a lymph node, more than 6 cm in greatest dimension

Distant Metastasis (pM)

- ☐ Not applicable
- ☐ pM1: Distant metastasis
Specify site(s), if known: _____