Melanoma Profile (Excision or Re-Excision)

Soft Code: C-MELEX

Procedure:	Excision, ellipse Excision, wide local Excision, other:		
Tumor Site/L	aterality:		
*Tumor Size: Tumor Site/S	cm Specimen laterality: Right Left Midline Not specified		
Tumor size (greatest dimension): cm Cannot be determined		
Histologic Ty	pe:		
Invasive:	No residual melanoma identified Superficial spreading Nodular Lentigo maligna Acral-lentiginous Desmoplastic Melanoma arising from blue nevus Melanoma arising in a giant congenital nevus Melanoma of childhood Nevoid melanoma Persistent melanoma Melanoma, not otherwise classified Other:		
In Situ (anato	Demic level 1) Superficial spreading Lentigo maligna type Acral-lentiginous Arising in a giant congenital nevus Melanoma in situ of childhood Persistent melanoma in situ Melanoma in situ, not otherwise classified Other:		

Revised February 2017 Soft Code: C-MELEX

Anatomic (Clark's) Level: At least level ____

II: Melanoma present in but does not fill and expand

papillary dermis

III: Melanoma fills and expands papillary dermis

IV: Melanoma invades reticular dermis

V: Melanoma invades subcutis

Maximum Tumor Thickness (Breslow depth): ____ mm

At least __ mm

Cannot be determined

Tumor-Infiltrating Lymphocytes:

Tumor Regression: Not identified

Present

Cannot be determined

Mitotic Rate (/mm2): None identified

≥1 mm² (specify number): _____

Growth Phase: Radial

Vertical

Cannot be determined

Microscopic Satellitosis: Not identified

Present

Cannot be determined

Macroscopic Satellitosis: Not identified

Present

Cannot be determined

Ulceration: Present

Not identified

Cannot be determined

Lympho-vascular Invasion: Not identified

Revised February 20	17		Soft Code: C-MELEX		
		Present Cannot be determined			
Perineural Invasion	:	Not identified Present Cannot be determined			
Margins:					
Peripheral M	largins: Canı	not be assessed			
	Unin	nvolved by invasive melanoma			
		Distance from closest margin: _	_ mm		
		Location:			
	Invo	lved by invasive melanoma; location	on:		
	Unin	nvolved by melanoma in situ			
		Distance from closest margin: _	_ mm		
		Location:			
	Invo	lved by melanoma in situ; location:	·		
Deep Margir	n: Canı	not be assessed			
, ,		nvolved by invasive melanoma			
		Distance from closest margin: _	mm		
	Invo	lved by invasive melanoma; location			
		•			
Co-existing Nevus:					
	Absent				
Lymph Nodes:	Total numb	per of nodes examined			
	Number of	sentinel lymph nodes examined:			
	Number of	lymph nodes with metastases:			
Pathologic Staging (pTNM):					
pTX: Primary tumor cannot be assessed (eg, shave biopsy or regressed melanoma)					
/aumlain)					

(explain): _____ pT0: No evidence of primary tumor

pTis: Melanoma in situ (ie, not an invasive tumor: anatomic level I) pT1: Melanoma 1.0 mm or less in thickness, with or without ulceration

pT1a: Melanoma 1.0 mm or less in thickness, no ulceration

pT1b: Melanoma 1.0 mm or less in thickness with ulceration and/or 1 or more mitoses/mm2

Soft Code: C-MELEX

pT2: Melanoma 1.01 to 2 mm in thickness, with or without ulceration

pT2a: Melanoma 1.01 to 2.0 mm in thickness, no ulceration

pT2b: Melanoma 1.01 to 2.0 mm in thickness, with ulceration

pT3: Melanoma 2.01 to 4.0 mm in thickness, with or without ulceration

pT3a: Melanoma 2.01 to 4.0 mm in thickness, no ulceration

pT3b: Melanoma 2.01 to 4.0 mm in thickness, with ulceration

pT4: Melanoma greater than 4.0 mm in thickness, with or without ulceration

pT4a: Melanoma greater than 4.0 mm in thickness, no ulceration

pT4b: Melanoma greater than 4.0 mm in thickness, with ulceration

^{*}The measurement of tumor size may underestimate the clinical size of the lesion due to tissue shrinkage