

# Gallbladder Carcinoma

Size: \_\_\_\_\_ cm

Site:

**Fundus**

**Body**

**Neck**

**Cystic duct**

**Free peritoneal side of gallbladder**

**Hepatic side of gallbladder**

**Other (specify): \_\_\_\_\_**

**Cannot be determined**

Type of carcinoma

**Ordinary adenocarcinoma**

**Other (specify): \_\_\_\_\_**

Histologic grade:

**Well differentiated**

**Moderately differentiated**

**Poorly differentiated**

**Undifferentiated**

**Other (specify): \_\_\_\_\_**

**Not applicable**

Extent of involvement:

**Lamina propria**

**Muscularis**

**Perimuscular connective tissue**

**Serosal surface (tumor is present on the surface)**

**Liver**

**Extrahepatic bile ducts**

**Adjacent organs (specify which organ or organs): \_\_\_\_\_**

**Main portal vein or hepatic artery**

Involvement of resection margins?: **Yes (specify which margin)** **No**

Positive lymph nodes/Total number of nodes: \_\_\_\_/\_\_\_\_

Distant metastasis: **Yes (specify)** **Unknown**

**pTNM**

## **T Stage**

TX primary tumor cannot be assessed

T1a invades lamina propria

T1b invades muscularis

T2 Tumor invades perimuscular connective tissue. No extension beyond the serosa or into the liver

T3 Tumor perforates the serosa (visceral peritoneum) and /or directly invades the liver or one other adjacent organ or structure such as the stomach, duodenum, colon, pancreas, omentum or extrahepatic bile ducts

T4 Tumor invades main portal vein or hepatic artery or invades 2 or more extrahepatic organs or structures

**N Stage**

NX Regional lymph nodes cannot be assessed

N0 No regional lymph node metastasis

N1 Metastasis to nodes along the cystic duct, common bile duct, hepatic artery, and/or portal vein

N2 Metastasis to periaortic, pericaval, superior mesenteric and /or celiac artery nodes

**DISTANT METASTASIS (M)**

M1 Distant metastasis (specify sites)

**If not M1, then do not include an M stage.**

**We cannot determine if there are no distant metastasis, so M0 cannot be used.**