THYROID CARCINOMA

Procedure:	Lobecto	omy					
	Total th	yroidectomy					
	Other:						
Tumor Focality:	Unifocal						
	Multifocal						
Tumor Site:	Right lobe						
	Left lobe						
	Isthmus						
	Pyramidal						
Tumor Size (greatest d	imensior	n): cm					
Histologic Type:	Papillary thyroid carcinoma, conventional type						
	Papillary thyroid carcinoma, follicular variant						
	Papillary thyroid carcinoma, tall cell variant						
	Papillary thyroid carcinoma, cribriform-morular variant						
	Papillary thyroid carcinoma, diffuse sclerosing variant						
Follicular carcinoma							
	Medulla	ary carcinoma					
	Poorly-differentiated carcinoma						
	Undifferentiated (anaplastic) carcinoma						
Other:							
Capsular Invasion:		Present	Absent		N/A		
Angioinvasion:		Present	Absent		N/A		
Lymphatic Invasion:		Present	Absent				
Extrathyroidal Extension:		Present (gros	ss)	Present (microscopic)			Absent
Margin Status:		Positive	Negativ	/e			

Extent of Primary Tumor:

- No evidence of primary tumor (pT0)
- o Tumor 1 cm or smaller in greatest dimension without extrathyroidal extension (pT1a)
- Tumor larger than 1 cm but not larger than 2 cm in greatest dimension without extrathyroidal extension (pT1b)
- Tumor larger than 2 cm but not larger than 4 cm in greatest dimension without extrathyroidal extension (pT2)
- o Tumor larger than 4 cm in greatest dimension without extrathyroidal extension (pT3a)
- Gross extrathyroidal extension invading only strap muscles (sternohyoid, sternothyroid, thyrohyoid, or omohyoid muscles) from a tumor of any size (pT3b)

- Gross extrathyroidal extension invading subcutaneous soft tissues, larynx, trachea, esophagus, or recurrent laryngeal nerve from a tumor of any size (pT4a)
- Gross extrathyroidal extension invading prevertebral fascia or encasing the carotid artery or mediastinal vessels from a tumor of any size (pT4b)

Lymph Node Status:

- Not assessed
- No regional lymph node metastasis (pN0)
- Metastasis to level VI or VII (pretracheal, paratracheal, prelaryngeal/Delphian, or upper mediastinal) lymph nodes (pN1a)
- Metastases to unilateral, bilateral, or contralateral lateral neck lymph nodes (levels I, II, III, IV, or V) or retropharyngeal lymph nodes (pN1b)

Distant Metastasis:

- Not assessed
- Distant metastasis (pM1)