

**URETHRA NEOPLASM**

Site: Male urethra  
       Penile  
       Bulbomembranous  
       Prostatic  
       Cannot be determined  
 Female urethra  
       Anterior  
       Posterior  
       Cannot be determine

Procedure: Cystoprostatectomy  
 Urethrectomy  
 Prostatectomy  
 Other: \_\_\_\_\_

Histologic Type: Urothelial carcinoma  
 Adenocarcinoma  
 Squamous cell carcinoma  
 Tumors of Mullerian type  
 Other: \_\_\_\_\_

Divergent Differentiation: Present Absent N/A

Grade (ISUP/WHO) Grade: Low-grade  
 High-grade  
 Well-differentiated for squamous cell carcinoma or adenocarcinoma  
 Moderately-differentiated for squamous cell carcinoma or adenocarcinoma  
 Poorly-differentiated for squamous cell carcinoma or adenocarcinoma  
 Other: \_\_\_\_\_

Architecture: Papillary Flat Inverted Mixed N/A

Size (maximum dimension): \_\_\_\_\_ cm

Extent of Primary Tumor (pT):

Primary tumor cannot be assessed (pTX)

No evidence of primary tumor (pT0)

Noninvasive papillary carcinoma (pTa)

Carcinoma in situ (pTis)

Tumor invades subepithelial connective tissue (pT1)

Tumor invades any of the following: corpus spongiosum, periurethral muscle (pT2)

Tumor invades any of the following: corpus cavernosum, anterior vagina (pT3)

Tumor invades other adjacent organs (pT4) (eg, invasion of the bladder wall)

For the Prostatic Urethra:

No evidence of primary tumor (pT0)

Non-invasive papillary carcinoma (pTa)

Carcinoma in situ involving the prostatic urethra or periurethral or prostatic ducts without stromal invasion (pTis)

Tumor invades urethral subepithelial connective tissue immediately underlying the urothelium (pT1)

Tumor invades the prostatic stroma surrounding ducts either by direct extension from the urothelial surface or by invasion from prostatic ducts (pT2)

Tumor invades the periprostatic fat (pT3)

Tumor invades other adjacent organs (pT4) (eg, invasion of the bladder wall)

Regional Lymph Nodes (pN):

Regional lymph nodes cannot be assessed (pNX)

No regional lymph node metastasis (pN0)

Single regional lymph node metastasis in the inguinal region or true pelvis (perivesical, obturator, internal (hypogastric) and external iliac) or presacral lymph node (pN1)

Multiple regional lymph node metastasis in the inguinal region or true pelvis (perivesical, obturator, internal (hypogastric) and external iliac) or presacral lymph node (pN2)

Number (fraction positive): \_\_\_\_\_ / \_\_\_\_\_

Site(s): \_\_\_\_\_

Size of largest metastatic deposit: \_\_\_\_\_ mm

Size of largest positive lymph node: \_\_\_\_\_ cm

Extranodal extension: Present Absent

Distant metastasis: N/A  
Present (pM1); site: \_\_\_\_\_

Adjacent Flat Carcinoma In Situ (CIS): Present Absent

Lymphovascular Invasion: Present Absent

Margin Status:

Distal urethral mucosal Negative  
Positive, noninvasive low-grade urothelial carcinoma/urothelial dysplasia  
Positive, CIS/noninvasive high-grade urothelial carcinoma  
Positive, invasive carcinoma

Proximal urethral mucosal Negative  
Positive, noninvasive low-grade urothelial carcinoma/urothelial dysplasia  
Positive, CIS/ noninvasive high-grade urothelial carcinoma  
Positive, invasive carcinoma  
NA

Soft tissue: Positive  
Negative