

## URINARY BLADDER NEOPLASM

Specimen: Urinary bladder and prostate  
 Urinary bladder  
 Other: \_\_\_\_\_

Procedure: Cystoprostatectomy  
 Cystectomy  
 Other: \_\_\_\_\_

Tumor Site: Anterior wall  
 Posterior wall  
 Dome  
 Right lateral wall  
 Left lateral wall  
 Trigone  
 Multifocal  
 Other: \_\_\_\_\_

Tumor Size (greatest dimension): \_\_\_\_\_ cm

Histologic Type: Urothelial carcinoma  
 Squamous cell carcinoma  
 Adenocarcinoma  
 Small cell carcinoma  
 Other: \_\_\_\_\_

Grade (ISUP/WHO): Low-grade  
 High-grade  
 Other: \_\_\_\_\_

Architecture: Papillary Inverted Mixed N/A

Non-urothelial Differentiation: Present Absent N/A  
 If present, specify: \_\_\_\_\_

Involvement of Prostate: Absent  
 Present, urethra only  
 Present, prostatic ducts/acini  
 Present, prostatic parenchyma  
 N/A

Adjacent Flat Carcinoma In Situ (Tis): Present Absent N/A

Lymphovascular Invasion: Present Absent

Margin Status:

Urethral:	Positive	Negative	N/A
Soft Tissue:	Positive	Negative	
Ureteral:	Positive	Negative	N/A

Extent of Primary Tumor:

- No evidence of primary tumor (pT0)
- Noninvasive papillary carcinoma (pTa)
- Flat carcinoma in situ (pTis)
- Tumor invades lamina propria (pT1)
- Tumor invades muscularis propria, inner half (pT2a)
- Tumor invades muscularis propria, outer half (pT2b)
- Tumor invades perivesicular tissue, microscopically (pT3a)
- Tumor invades perivesicular tissue, macroscopically (pT3b)
- Tumor invades prostate parenchyma/uterus/vagina\* (pT4a)
- Tumor invades pelvic wall/abdominal wall (pT4b)

*\*Defined as prostatic stromal invasion directly from bladder cancer. This staging system should not be used for tumors originating in prostatic urethra. Subepithelial invasion of prostatic urethra of a tumor from the prostatic urethra does not constitute T4 staging status.*

Lymph Node Status:

- No lymph node metastasis (pN0)
- Single regional lymph node metastasis in the true pelvis (hypogastric, obturator, external iliac or presacral lymph node) (pN1)
- Metastasis in multiple regional lymph nodes in the true pelvis (pN2)
- Lymph node metastasis to the common iliac lymph nodes (pN3)

Number (fraction positive): \_\_\_\_\_ / \_\_\_\_\_

Site(s): \_\_\_\_\_

Size of largest positive lymph node: \_\_\_\_\_ cm

Extranodal extension: Present                  Absent